

ONSET OF CAVITY DEFORMATION UPON SUBSONIC MOTION OF A PROJECTILE IN A FLUID COMPLEX PLASMA

D. I. Zhukhovitskii,^{1,*} A. V. Ivlev,² V. E. Fortov,¹ and G. E. Morfill²

¹*Joint Institute of High Temperatures, Russian Academy of Sciences,*

Izhorskaya 13, Bd. 2, 125412 Moscow, Russia

²*Max-Planck-Institut für extraterrestrische Physik,*

Giessenbachstrasse, 85748 Garching, Germany

(Dated: January 29, 2013)

We study deformation of a cavity around a large projectile moving with subsonic velocity in the cloud of small dust particles. To solve this problem, we employ the Navier-Stokes equation for a compressible fluid with due regard for friction between dust particles and atoms of neutral gas. The solutions shows that due to friction, the pressure of dust cloud at the boundary of the cavity behind the projectile can become negative, which entails formation of a microscopic void free from dust particles – the cavity deformation. The corresponding threshold velocity is calculated, which is found to decrease with increasing cavity size. Measurement of such velocity makes it possible to estimate the static pressure inside the dist cloud.

PACS numbers: 52.27.Lw, 83.10.Rs, 82.70.Dd

* dmr@ihed.ras.ru

I. INTRODUCTION

Complex (dusty) plasma is a low-temperature plasma including microparticles. Due to electron absorption, particles acquire a considerable electric charge. Thus, a strongly coupled Coulomb system is formed [1–9]. Such plasma represents a natural system which makes it possible to observe various collective phenomena at the level of individual particles. In experimental setups, complex plasmas are usually studied in gas discharges at low pressures, e.g., in radio frequency (RF) discharges. A large homogeneous bulk of complex plasma, which almost fills the entire discharge volume, can be observed under microgravity conditions either in parabolic flights [10–14] or onboard the International Space Station (ISS) [10, 15–19].

In recent studies, attention was focused on tracer particles or projectiles moving through a cloud of complex plasma. Such projectiles are generated using controlled mechanisms of acceleration [11, 20], or they can appear sporadically [19, 21]. Projectiles moving with supersonic velocity lead to the formation of extended Mach cones, subsonic (slow) ones produce localized disturbances of surrounding particles. In Ref. 22, it was suggested that the latter regime, realized when a relatively large subsonic projectile moves in a dense cloud of smaller particles, can be well approximated by a flow of an incompressible fluid. In the framework of the same hydrodynamic approximation, it was demonstrated that, along with the regular neutral gas drag, there is an additional force exerted on the projectile due to friction between neutral atoms and the particle fluid [23].

In this study, we try to estimate the deformation threshold for a (initially spherical) cavity around a subsonic projectile. This deformation emerges abruptly as the projectile velocity increases, as it is seen in snapshots given in [11, 22]. To solve this problem, we employ

the Navier-Stokes equation for a compressible fluid taking into account friction between dust particles and atoms of neutral gas. We do not treat the deformation self-consistently. Instead, we imply that the projectile velocity is below the threshold value, so that a regular flow around a spherical cavity with no stall can be treated. The solutions shows that due to friction, the pressure of dust cloud at the boundary of the cavity behind the projectile can become negative, which entails the formation of a microscopic void free from dust particles – the cavity deformation. This occurs at some threshold velocity which decreases with increasing cavity size. Measurement of such velocity would make it possible to estimate the static pressure inside the dust cloud.

The paper is organized as follows. In Sec. II, we solve the non-stationary Navier-Stokes equation for an incompressible particle fluid, in Sec. III, the gas dynamics problem is solved and the corrections for fluid compressibility to the velocity and pressure fields are calculated. Calculation results are compared with available experimental data in Sec. IV, the results of this study are summarized in Sec. V.

II. INCOMPRESSIBLE FLUID APPROXIMATION

Consider an irrotational flow of incompressible particle fluid formed by the dust crystal melted around a projectile moving with the velocity $\mathbf{u}(t)$ relative to the dust, where t is the time. In this Section, we will assume that the cavity around a projectile is not deformed and has the form of a sphere with radius R . Note that this is the only possible form of a cavity when the parameter of interaction between the projectile and dust particles, β_{dp} [22], is great (under typical experimental conditions, $\beta_{dp} \sim 100$ [22]). The Navier-Stokes equation

describing the velocity field $\mathbf{v}(\mathbf{r}, t)$ in the reference frame of the projectile has the form

$$\frac{\partial \mathbf{v}}{\partial t} + (\mathbf{v} \cdot \nabla) \mathbf{v} + \nu(\mathbf{v} + \mathbf{u}) = -\frac{\nabla p}{\rho}, \quad (1)$$

where $\rho = M_d n_d$ is the mass density of dust fluid (assumed in this section to be constant), with M_d and n_d being, respectively, the mass and number density of dust particles of the radius a_d . Furthermore, $p(t, \mathbf{r})$ is the dust pressure field, $\nu = (8\sqrt{2\pi}/3)\delta m_n n_n v_{Tn} a_d^2 / M_d$ is the friction coefficient with $\delta \simeq 1.4$ being the accommodation coefficient [24], and m_n , n_n , T_n , and $v_{Tn} = (T_n/m_n)^{1/2}$ are the mass, number density, temperature, and thermal velocity of neutral gas molecules, respectively. Equation (1) assumes also that far from the projectile, dust particles are quiescent relative to neutral gas. It was shown in [22] that the approximation of nonviscous flow results in a fairly good description of trajectories of individual dust particles; estimate of the viscosity term for the dust particle fluid is indicative of the fact that in most cases, it is small [23]. This allowed us to omit it in Eq. (1) and to confine ourselves to the nonviscous approximation.

For an incompressible fluid, the continuity equation is reduced to

$$\nabla \cdot \mathbf{v} = 0. \quad (2)$$

The boundary conditions for (1) and (2) are $(\mathbf{v} \cdot \mathbf{n})|_{r=R} = 0$, where $\mathbf{n} = \mathbf{r}/r$, and $\mathbf{v} = -\mathbf{u}$ at $r = \infty$. For an irrotational flow ($\nabla \times \mathbf{v} = 0$), we substitute $\mathbf{v} = \nabla \varphi - \mathbf{u}$ in Eq. (2) to obtain the equation

$$\nabla^2 \varphi = 0, \quad \varphi(t, \infty) = 0, \quad \left. \frac{\partial \varphi}{\partial \mathbf{n}} \right|_{r=R} = 0, \quad (3)$$

which has the solution [25]

$$\varphi(\mathbf{r}) = -\frac{R^3}{2r^2} \mathbf{u} \cdot \mathbf{n} \quad (4)$$

or

$$\mathbf{v}(\mathbf{r}) = -\frac{R^3}{2r^3} [\mathbf{u} - 3\mathbf{n}(\mathbf{n} \cdot \mathbf{u})] - \mathbf{u}. \quad (5)$$

The pressure field $p(\mathbf{r})$ can then be found by substituting velocity (5) in Eq. (1). This yields the following pressure distribution at the spherical surface of a cavity ($r = R$),

$$p = p_{\text{st}} + \frac{\rho\nu R}{2}\mathbf{u} \cdot \mathbf{n} + \frac{\rho u^2}{8} \left[9 \frac{(\mathbf{u} \cdot \mathbf{n})^2}{u^2} - 5 \right] + \frac{\rho R}{2} \dot{\mathbf{u}} \cdot \mathbf{n}, \quad (6)$$

where $p_{\text{st}} = \text{const} > 0$ is the static pressure of dust at $r = \infty$.

Consider deformation of a cavity around a projectile propagating in the dust crystal (Fig. 1). Here, we will neglect finite fluid compressibility – corresponding corrections are discussed in Sec. III. Deformation can be caused solely by formation of a void in the stream of dust fluid because the interaction parameter β_{dp} is large. Since no evidence for an appreciable attraction between dust particles has been reported in the literature, we model the cloud of dust particles by a system of repulsing soft spheres. In such a system, a cavity can be deformed by void formation if the condition $p \leq 0$ is satisfied over some area around a cavity. This condition corresponds to the cavitation condition in a metastable fluid and to stall in gas dynamics. One can associate the shape of a cavity with the line $p(\mathbf{r}) = 0$.

Let us first consider the case $\dot{u} = 0$. According to Eq. (6), friction between dust particles and neutral gas gives rise to pressure increase in front of the projectile and its decrease behind it, so that $p(\mathbf{r})$ reaches a maximum at the “front pole” (where $\mathbf{u} \cdot \mathbf{n} = u$). By introducing the cosine of the polar angle, $\cos \theta = \mathbf{u} \cdot \mathbf{n}/u \equiv \xi$, we readily derive

$$\frac{\partial p}{\partial \xi} = \frac{9}{4} \rho u^2 (\xi - \xi_{\text{cr}}), \quad (7)$$

where $\xi_{\text{cr}} = -\frac{2}{9}(\nu R/u)$. If $|\xi_{\text{cr}}| > 1$ then the pressure reaches a minimum at the “rear pole” ($\xi = -1$), where $p_{\text{min}} = p_{\text{st}} - \frac{1}{2} \rho u^2 (\frac{9}{2} |\xi_{\text{cr}}| - 1)$. This corresponds to the low-velocity regime $u < \frac{2}{9} \nu R$, when the minimum pressure decreases (approximately) linearly both with u and R . The cavity deformation threshold is defined by the condition $p_{\text{min}} = 0$ and is reached easier for larger projectiles, at the critical velocity $u_{\text{cr}} \simeq 2p_{\text{st}}/(\rho\nu R)$ [here

we assume $p_{\text{st}} \lesssim \frac{1}{8}\rho(\nu R)^2$. Note that in the high-velocity regime $u > \frac{2}{9}\nu R$ the pressure minimum is shifted to a certain latitude in the rear hemisphere, approaching the “equator” ($\xi = 0$) asymptotically. In this case the friction plays minor role and the critical velocity is $u_{\text{cr}} \simeq \sqrt{8p_{\text{st}}/5\rho}$.

It is well known in hydrodynamics that the presence of points or small regions of negative pressure can lead to cavitation. However, in complex plasmas the distance between fluid particles is likely to be even greater than the size of such regions. Therefore, the continuity approximation fails to describe the cavitation in complex plasmas and, hence, this process is unlikely to be observed in a real experiment. On the other hand, upon further increase of the projectile velocity the regions of negative pressure should grow until they cover an appreciable part of the surface of a cavity, including its rear pole. This should be followed by a stall observed as cavity deformation. At this stage, a growing deviation from the spherical geometry makes analytical solution of the problem too complicated. Moreover, to our best knowledge, there is no criterion that would tell us how large the negative region should be to initiate a stall. Nevertheless, one can speculate that the velocity u_{cr} derived above provides a reasonable order-of-magnitude estimate for the cavity deformation onset, so that one can also estimate the static pressure p_{st} .

Note that for arbitrary acceleration (assuming $\dot{\mathbf{u}}\|\mathbf{u}$), one can introduce the effective damping rate $\nu_{\text{eff}} = \nu + \dot{u}/u$ which can have either sign, so that the critical cosine ξ_{cr} can have either sign as well. Thus, if a projectile experiences sufficiently strong deceleration ($|\dot{u}|/u > \nu$) the cavity can be also deformed at the front hemisphere.

III. FLOW WITH FINITE COMPRESSIBILITY

In this Section, we will estimate the effect of finite compressibility of the dust cloud on the velocity and pressure fields. Here, we will treat a steady ($\dot{\mathbf{u}} = 0$) and irrotational ($\nabla \times \mathbf{v} = 0$) flow, so that the fluid density is a function solely of the radius-vector: $\rho = \rho(\mathbf{r})$. Since $\nabla \times \mathbf{v} = 0$ and $\nabla p = c^2 \nabla \rho$, where $c = \sqrt{(\partial p / \partial \rho)_T}$ is the sound velocity, we can rewrite Eqs. (1) and (2) in the form

$$(\mathbf{v} \cdot \nabla) \mathbf{v} + \nu(\mathbf{v} + \mathbf{u}) = -\frac{c^2}{\rho} \nabla \rho. \quad (8)$$

In this case, the continuity equation includes the fluid density,

$$\nabla \cdot \mathbf{v} + \mathbf{v} \cdot \frac{\nabla \rho}{\rho} = 0. \quad (9)$$

Here, c is assumed to be constant independent of the local velocity and plasma state parameters. We substitute $\nabla \rho / \rho = -(1/c^2)(\mathbf{v} \cdot \nabla) \mathbf{v} - (\nu/c^2)(\mathbf{v} + \mathbf{u})$ from (8) in (9) to derive

$$\nabla \cdot \mathbf{v} = \frac{1}{c^2} \mathbf{v} \cdot (\mathbf{v} \cdot \nabla) \mathbf{v} + \frac{\nu}{c^2} (v^2 + \mathbf{u} \cdot \mathbf{v}). \quad (10)$$

We will search for the solution of Eq. (10) in the form $\mathbf{v} = \mathbf{v}_0 + \mathbf{v}_1$, where $\mathbf{v}_0(\mathbf{r})$ is the solution of incompressible problem (for $\rho = \rho_0 = \text{const}$, i.e., for $c \rightarrow \infty$) given by Eq. (5), the respective pressure distribution $p_0(\mathbf{r})$ is presented by Eq. (6) (for $\dot{u} = 0$). Obviously, $\nabla \cdot \mathbf{v}_0 = 0$. We assume that $|\mathbf{v}_1| \ll |\mathbf{v}_0|$ and retain solely zeroth-order terms on the rhs of (10) to obtain

$$\nabla \cdot \mathbf{v}_1 = \frac{1}{2c^2} \mathbf{v}_0 \cdot \nabla v_0^2 + \frac{\nu}{c^2} (v_0^2 + \mathbf{u} \cdot \mathbf{v}_0). \quad (11)$$

Here, we took into account that $\nabla \times \mathbf{v}_0 = 0$ and hence $(\mathbf{v}_0 \cdot \nabla) \mathbf{v}_0 = \nabla v_0^2 / 2$. Since $|\mathbf{v}_0| \sim u$, $|\nabla| \sim 1/R$, and $(u/c)^2$ is small for subsonic motion, the estimate $|\mathbf{v}_1| \sim (u^2/c^2) \max\{u, \nu R\}$ following from (11) justifies our assumption. It is seen from this estimate that both sides

of Eq. (11) are of the order $(u/c)^2$, and terms other than zeroth-order ones would lead to higher-order terms in this parameter. Obviously, $|\mathbf{v}_1| \rightarrow 0$ at $c \rightarrow \infty$.

We introduce the scaled coordinate $x = r/R$ and the dimensionless potential $\tilde{\varphi}_1$ defined by the relation $\mathbf{v}_1 = (u^3/c^2)\nabla\tilde{\varphi}_1$ (in so doing, the condition $\nabla \times \mathbf{v}_1 = 0$ is satisfied) to rewrite (11) in the dimensionless form

$$\begin{aligned} \nabla^2 \tilde{\varphi}_1 &= \frac{\kappa}{2x^6} P_0(\xi) + \left(\frac{36}{5x^7} - \frac{9}{2x^{10}} \right) P_1(\xi) \\ &- \kappa \left(\frac{1}{x^3} - \frac{1}{2x^6} \right) P_2(\xi) - \left(\frac{6}{x^4} - \frac{24}{5x^7} + \frac{3}{2x^{10}} \right) P_3(\xi), \end{aligned} \quad (12)$$

where $P_l(\xi)$ are the Legendre polynomials and $\kappa = \nu R/u$. Here and in what follows, differentiation with respect to scaled coordinates is assumed. The boundary conditions for Eq. (12) follow from the conditions $\mathbf{v}_1(\infty) = 0$ and $\mathbf{n} \cdot \mathbf{v}_1|_{r=R} = 0$, i.e.,

$$\tilde{\varphi}_1(\infty) = 0, \quad \left. \frac{d\tilde{\varphi}_1}{d\mathbf{n}} \right|_{x=1} = 0. \quad (13)$$

Solution of Eq. (12) is derived in Appendix A. The resulting finite-compressibility corrections to the potential and velocity *at the cavity surface* follow from (A6):

$$\begin{aligned} \tilde{\varphi}_1|_{x=1} &= \frac{81}{176}\xi^3 + \frac{\kappa}{2} \left(\frac{1}{4}\xi^2 - \frac{1}{3} \right) - \frac{551}{880}, \\ \mathbf{v}_1|_{x=1} &= \frac{u^2}{c^2} \left(\frac{243}{176}\xi^2 + \frac{\kappa}{4}\xi - \frac{551}{880} \right) (\mathbf{u} - \xi \mathbf{un}). \end{aligned} \quad (14)$$

As it could be expected from symmetry considerations, $\mathbf{v}_1|_{x=1} = 0$ at $\xi = \pm 1$, i.e., at both poles. Note that result (14) validates the estimate $|\mathbf{v}_1| \sim (u^2/c^2) \max\{u, \nu R\}$ given above.

On the basis of velocity correction (14) and using Eq. (8), it is straightforward to calculate a compressibility correction to the pressure, $p_1 = p - p_0$. This is performed in Appendix B. It is worth mentioning that in the general case, correction to the pressure (B4) includes solely even powers of ξ , so that corresponding correction to an additional force exerted on the projectile due to dissipation in the surrounding particle fluid [Eq. (10) in Ref. [23]]

vanishes. In the friction-dominated regime (large κ), one can retain in (B4) solely the term proportional to u^2 , which is independent of ξ , so that the total pressure is

$$p(\xi) \simeq p_{\text{st}} + \frac{\rho_0 u \nu R}{2} \xi + \frac{\rho_0 u^2}{8} \left(9\xi^2 - 5 + \frac{4\nu^2 R^2}{3c^2} \right). \quad (15)$$

Thus, the pressure increases due to finite compressibility, so that the threshold velocity of cavity deformation increases as well. On the other hand, one can see that the resulting correction is small when $\kappa(u/c)^2 \ll 1$. As we show in the next section, the compressibility correction can usually be neglected for subsonic projectiles under typical experimental conditions.

IV. COMPARISON WITH EXPERIMENT

Based on the developed theory, let us analyze the possibility of cavity deformation in typical experiments with complex plasmas. We first focus on the experiments carried out in the PK-3 Plus laboratory onboard the ISS under microgravity conditions. Details on the setup can be found in [18]. Dust particles injected into the main plasma formed a cloud around the center of the chamber. Several larger particles present in the chamber get sporadically accelerated and penetrate the cloud, thus generating projectiles [26]. Consider the experiment performed with argon gas ($m_n = 6.6 \times 10^{-23}$ g). For the gas pressure 10 Pa and temperature $T_n = 300$ K we have $n_n = 2.4 \times 10^{15} \text{ cm}^{-3}$ and $v_{T_n} = 2.5 \times 10^4 \text{ cm/s}$. Dust cloud was formed by melamine-formaldehyde particles with the diameter $2a_d = 2.55 \times 10^{-4} \text{ cm}$ and mass $M_d = 1.3 \times 10^{-11} \text{ g}$ [19, 22], therefore, $\nu \simeq 46 \text{ s}^{-1}$. The path of a subsonic projectile ($u < c \simeq 2.1 \text{ cm/s}$ [19], recorded with a high-resolution camera) revealed cavity deformation. The most obvious manifestation of the deformation is a visible shift of the projectile from cavity center to its front side. This shift can be accompanied by cavity

elongation (Fig. 2). In the course of motion of the projectile, its velocity slowly increased from $u \simeq 0.7$ cm/s to 1.4 cm/s. Cavity deformation becomes visible approximately in the middle of the path; the projectile velocity and acceleration at this point are 1.06 cm/s and 2.6 cm/s², respectively, the radius of a spherical cavity around the projectile at this point is $R \simeq 3.7 \times 10^{-2}$ cm.

One can crudely estimate the static pressure p_{st} of dust cloud from the observed deformation threshold. According to the discussion at the end of Sec. II, the projectile velocity at which a point of negative pressure emerges on the cavity surface, is lower than the velocity of deformation onset observed experimentally. Hence, in this case $2\nu R/9u_{\text{cr}} \sim 1$, and the system is at the boundary between the low- and high-velocity regimes. It is not surprising therefore that the corresponding estimates for the static pressure, $K(\equiv p_{\text{st}}/n_d) \simeq M_d \nu R u_{\text{cr}}/2$ and $K \simeq 5M_d u_{\text{cr}}^2/8$, both yield $K \sim 3$ eV, i.e., $p_{\text{st}} \sim 10^{-7}$ Pa. Note that the correction for finite compressibility, $\nu^2 R^2/3c^2$, does not exceed 20%, and the correction for acceleration is negligible.

Microgravity conditions can also be maintained on parabolic flights. Such experiments were performed using the IMPF-K2 chamber [11]. The experiment was carried out with argon at 30 Pa and $2a_d = 9.55 \times 10^{-4}$ cm ($M_d = 6.8 \times 10^{-10}$ g and $\nu \simeq 33$ s⁻¹). Projectiles were accelerated by a special device up to supersonic or subsonic velocities. For a subsonic projectile, the cavity radius could be roughly estimated as $R \simeq 0.1$ cm. The projectile velocity amounts to $u \simeq 1.8$ cm/s, and cavity deformation is clearly seen in Fig. 3a of Ref. [11]. Since we have no information concerning the dependence of K on the size of a dust particle, we use the same value of K as for the experiment of Ref. [22] discussed above. Then the threshold velocity for this case does not exceed 1 cm/s. This corroborates the main notion of this study – the cavity around a moving projectile is normally deformed at

subsonic velocities.

V. CONCLUSION

We have investigated the possibility of deformation of a cavity around large projectiles slowly moving inside a cloud of small dust particles. This cavity is formed due to a strong Coulomb repulsion between the projectile and dust particles. Since the parameter β_{dp} characterizing this interaction is very large, cavity deformation occurs solely due to formation of a void adjacent to initially spherical cavity. This is similar to the process of void formation in a fluid under negative pressure. Since we approximate the dust cloud by a system of soft spheres, zero pressure is sufficient for void formation leading to cavity deformation.

We model a collective particle subsonic motion in a complex plasma by nonviscous irrotational hydrodynamic motion of dust fluid and show that, in this model, the reason for cavity deformation is the friction between dust particles and neutral atoms of a carrier gas. It is this friction that stipulates emergence of a zone of negative pressure over the back of a cavity surface. If such zone covers substantial part of the surface, stall takes place, which is observed as cavity deformation. We have found that the deformation is more likely to be observed for larger projectiles. The deformation occurs when the projectile velocity exceeds some threshold subsonic velocity.

Since typical projectile velocities are of the same order of magnitude as the sound velocity, we have studied the effect of finite compressibility of the dust fluid on velocity and pressure fields. With this purpose, we have obtained an exact solution of linearized gas dynamics equations for compressible nonviscous fluid flowing about the cavity. In so doing, we included both the convective and friction terms in the Navier-Stokes equation. Calculations showed that corrections for finite compressibility are rather small. Generally, account of the

compressibility leads to a slight shift of the area covered by a negative pressure zone.

Analysis of available experimental data has validated our approach. It was found that the cavity is deformed when either the velocity [22] or the size [11] of a projectile are sufficiently large. It is worth mentioning that the phenomenon of cavity deformation is similar to the formation of a void behind a cluster of smaller particles moving in the cloud of larger ones [27]. For intermediate projectile sizes, a transition from spherical to deformed cavity was observed along the trajectory of an individual projectile. The transition threshold is in accordance with the developed theory. This made it possible to estimate the static pressure inside a dust cloud and, hence, opened up a possibility to use projectiles as a diagnostic tool for the dust equation of state.

In this study, we restricted ourselves to investigation of the threshold of cavity deformation and did not treat the shape of deformed cavity as well as displacement of the projectile away from the center of its cavity. This problem, besides all, would require an adequate account for local non-spherical charge distribution around the projectile. It is also possible that cavity deformation promotes the observed shift of the projectile from the center of a cavity, driven by an additional drag force exerted on the projectile [23].

Appendix A:

The solution of Eq. (12) with boundary conditions (13) is represented in the form

$$\tilde{\varphi}_1(x, \xi) = \sum_{l=0}^3 f_l(x) P_l(\xi), \quad (\text{A1})$$

where $f_l(x)$ are the coefficients of expansion in the Legendre polynomials $P_l(\xi)$. We substitute (A1) in (12) taking into account that $\nabla^2 P_l(\xi) = -l(l+1)x^{-2}P_l(\xi)$ to reduce Eq. (12) to

a set of ordinary differential equations in spherical coordinates for nonzero functions $f_l(x)$,

$$\frac{d}{dx} \left(x^2 \frac{df_l}{dx} \right) - l(l+1)f_l = q_l(x), \quad (\text{A2})$$

where $q_0(x) = \frac{1}{2}\kappa x^{-4}$, $q_1(x) = \frac{36}{5}x^{-5} - \frac{9}{2}x^{-8}$, $q_2(x) = -\kappa x^{-1} + \frac{1}{2}\kappa x^{-4}$, $q_3(x) = -6x^{-2} + \frac{24}{5}x^{-5} - \frac{3}{2}x^{-8}$. The boundary conditions for (A2) follow from (13):

$$f_l(\infty) = 0, \quad \left. \frac{df_l}{dx} \right|_{x=1} = 0. \quad (\text{A3})$$

The rhs of (A2) has the form $\sum_k b_{lk}x^{-k}$, where b_{lk} are constants entering the functions $q_l(x)$.

Solution of (A2) is then given by

$$f_l(x) = \frac{C}{x^{l+1}} + \sum_k \frac{b_{lk}}{k(k-1) - l(l+1)} \frac{1}{x^k}, \quad (\text{A4})$$

where C is defined by the second boundary condition (A3), whence it follows that

$$f_l(x) = \sum_k \frac{b_{lk}}{l(l+1) - k(k-1)} \left(\frac{k}{l+1} \frac{1}{x^{l+1}} - \frac{1}{x^k} \right). \quad (\text{A5})$$

Finally, we obtain

$$\begin{aligned} f_0 &= \kappa \left(-\frac{1}{6x} + \frac{1}{24x^4} \right), \\ f_1 &= -\frac{2}{3x^2} + \frac{2}{5x^5} - \frac{1}{12x^8}, \\ f_2 &= \kappa \left(\frac{1}{6x} - \frac{1}{6x^3} + \frac{1}{12x^4} \right), \\ f_3 &= \frac{3}{5x^2} - \frac{54}{55x^4} + \frac{3}{5x^5} - \frac{3}{88x^8}. \end{aligned} \quad (\text{A6})$$

Appendix B:

The pressure correction due to finite compressibility can be found using the corresponding velocity correction, Eq. (14). We rewrite Eq. (8) in the form

$$-c^2 \nabla \ln \rho = \frac{1}{2} \nabla v^2 + \nu \nabla \varphi, \quad (\text{B1})$$

where $\varphi = \varphi_0 + \varphi_1$ and $\mathbf{v} = \mathbf{u} + \nabla\varphi$. By integrating Eq. (B1) and using Bernoulli's equation for compressible fluids we derive

$$\rho - \rho_0 = -\frac{\rho_0}{c^2} \left(\frac{v_0^2}{2} + \nu\varphi_0 \right) - \frac{\rho_0}{c^2} (\mathbf{v}_0 \cdot \mathbf{v}_1 + \nu\varphi_1) + \frac{\rho_0}{2c^4} \left(\frac{v_0^2}{2} + \nu\varphi_0 \right)^2, \quad (\text{B2})$$

where $\rho_0 = \rho|_{r=\infty}$ and the terms up to $(u/c)^4$ are retained (for simplicity we assume $c = \text{const}$). We write the equation of state for the dust fluid in the form $p - p_{\text{st}} = c^2(\rho - \rho_0)$, which yields

$$p = p_{\text{st}} - \frac{\rho_0 v_0^2}{2} - \nu\rho_0\varphi_0 - \rho_0\mathbf{v}_0 \cdot \mathbf{v}_1 - \nu\rho_0\varphi_1 + \frac{1}{2\rho_0 c^2} \left(\frac{\rho_0 v_0^2}{2} + \nu\rho_0\varphi_0 \right)^2. \quad (\text{B3})$$

The first three terms of Eq. (B3) represent Eq. (6) with $\dot{u} = 0$. Hence, by substituting in Eq. (B3) the expressions for φ_0 and $\mathbf{v}_0$ [Eqs (4) and (5)] as well as for φ_1 and $\mathbf{v}_1$ [Eq. (14)] taken at the cavity surface, we get

$$p_1(\xi) = \frac{\rho_0 u^4}{c^2} \left[\frac{1}{6} \kappa^2 - 3\kappa\xi^2 \left(\xi^2 - \frac{101}{440} \right) - \frac{2025}{1408} \left(\xi^4 - \frac{5414}{3375} \xi^2 + \frac{5237}{10125} \right) \right]. \quad (\text{B4})$$

-
- [1] V. E. Fortov and G. E. Morfill, eds., *Complex and Dusty Plasmas: From Laboratory to Space*, Series in Plasma Physics (CRC Press, 2009).
- [2] J. H. Chu and L. I., Phys. Rev. Lett. **72**, 4009 (1994).
- [3] H. Thomas, G. E. Morfill, V. Demmel, J. Goree, B. Feuerbacher, and D. Möhlmann, Phys. Rev. Lett. **73**, 652 (1994).
- [4] Y. Hayashi and S. Tashibana, Jpn. J. Appl. Phys. **33**, L804 (1994).
- [5] S. V. Vladimirov, K. Ostrikov, and A. A. Samarian, *Physics and applications of complex plasmas* (Imperial College, London, 2005).
- [6] V. Fortov, A. Ivlev, S. Khrapak, A. Khrapak, and G. Morfill, Phys. Rep. **421**, 1 (2005).
- [7] G. E. Morfill and A. V. Ivlev, Rev. Mod. Phys. **81**, 1353 (2009).
- [8] P. K. Shukla and B. Eliasson, Rev. Mod. Phys. **81**, 25 (2009).
- [9] M. Bonitz, C. Henning, and D. Block, Rep. Prog. Phys. **73**, 066501 (2010).
- [10] G. E. Morfill, U. Konopka, M. Kretschmer, M. Rubin-Zuzic, H. M. Thomas, S. K. Zhdanov, and V. Tsytovich, New J. Phys. **8**, 7 (2006).
- [11] D. Caliebe, O. Arp, and A. Piel, Phys. of Plasmas **18**, 073702 (2011).
- [12] A. Piel, O. Arp, M. Klindworth, and A. Melzer, Phys. Rev. E **77**, 026407 (2008).
- [13] K. O. Menzel, O. Arp, and A. Piel, Phys. Rev. E **83**, 016402 (2011).
- [14] O. Arp, D. Caliebe, and A. Piel, Phys. Rev. E **83**, 066404 (2011).
- [15] M. Schwabe, S. K. Zhdanov, H. M. Thomas, A. V. Ivlev, M. Rubin-Zuzic, G. E. Morfill, V. I. Molotkov, A. M. Lipaev, V. E. Fortov, and T. Reiter, New J. Phys. **10**, 033037 (2008).

- [16] G. E. Morfill, H. M. Thomas, U. Konopka, H. Rothermel, M. Zuzic, A. Ivlev, and J. Goree, Phys. Rev. Lett. **83**, 1598 (1999).
- [17] S. A. Khrapak, B. A. Klumov, P. Huber, V. I. Molotkov, A. M. Lipaev, V. N. Naumkin, H. M. Thomas, A. V. Ivlev, G. E. Morfill, O. F. Petrov, V. E. Fortov, Y. Malentschenko, and S. Volkov, Phys. Rev. Lett. **106**, 205001 (2011).
- [18] H. M. Thomas, G. E. Morfill, V. E. Fortov, A. V. Ivlev, V. I. Molotkov, A. M. Lipaev, T. Hagl, H. Rothermel, S. A. Khrapak, R. K. Suetterlin, M. Rubin-Zuzic, O. F. Petrov, V. I. Tokarev, and S. K. Krikalev, New J. Phys. **10**, 033036 (2008).
- [19] M. Schwabe, K. Jiang, S. Zhdanov, T. Hagl, P. Huber, A. V. Ivlev, A. M. Lipaev, V. I. Molotkov, V. N. Naumkin, K. R. Sütterlin, H. M. Thomas, V. E. Fortov, G. E. Morfill, A. Skvortsov, and S. Volkov, EPL **96**, 55001 (2011).
- [20] M.-C. Chang, Y.-P. Tseng, and L. I, Phys. of Plasmas **18**, 033704 (2011).
- [21] D. Samsonov, J. Goree, H. M. Thomas, and G. E. Morfill, Phys. Rev. E **61**, 5557 (2000).
- [22] D. I. Zhukhovitskii, V. E. Fortov, V. I. Molotkov, A. M. Lipaev, V. N. Naumkin, H. M. Thomas, A. V. Ivlev, M. Schwabe, and G. E. Morfill, Phys. Rev. E **86**, 016401 (2012).
- [23] A. V. Ivlev and D. I. Zhukhovitskii, Phys. Plasmas **19**, 093703 (2012).
- [24] V. Fortov, A. Ivlev, S. Khrapak, A. Khrapak, and G. Morfill, Phys. Rep. **421**, 1 (2005).
- [25] L. D. Landau and E. M. Lifshits, *Fluid Mechanics* (Pergamon Press, 1959).
- [26] J. Goree, G. E. Morfill, V. N. Tsytovich, and S. V. Vladimirov, Phys. Rev. E **59**, 7055 (1999).
- [27] K. R. Sütterlin, A. Wysocki, A. V. Ivlev, C. R  th, H. M. Thomas, M. Rubin-Zuzic, W. J. Goedheer, V. E. Fortov, A. M. Lipaev, V. I. Molotkov, O. F. Petrov, G. E. Morfill, and H. L  wen, Phys. Rev. Lett. **102**, 085003 (2009).

FIGURE CAPTIONS

Fig. 1: Deformation of a spherical cavity (of the initial radius R) behind a projectile (large bullet in the center). The projectile moves with the velocity u through a dust cloud (small bullets around) embedded in a carrier gas (gray background).

Fig. 2: Almost spherical (a) and deformed (b) cavity around a projectile at different portions of its trajectory, observed with a high-resolution camera (inverted image).




FIG. 1.




FIG. 2.